

1994 Guidelines	2013 Guideline
<p style="text-align: center;">Behavior/History Exclusionary Criteria</p> <p><i>Specific Exclusionary Criteria for Ped Donors #1-</i> (Children meeting any of the exclusionary criteria listed below for adults should not be accepted as donors)</p>	<p><i>Donors who meet one or more of the following 11 criteria should be identified as being at increased risk for recent HIV, HBV, and HCV infections. Each factor listed reflects increased risk of all 3 pathogens as an aggregate, as there is overlap of associated risk, even though each factor does not convey risk from all pathogens equally. The first six factors address sexual contact; the definition of "had sex" refers to any method of sexual contact, including vaginal, anal and oral contact:</i></p>
#5- Persons who have had sex in the preceding 12 months with any person described in items 1-4 or with a person known or suspected to have HIV infection	People who have had sex with a person known or suspected to have HIV, HBV, or HCV infections in the preceding 12 months
#1- Men who have had sex with another man in the preceding 5 years	MSM in the preceding 12 months
#5- Persons who have had sex in the preceding 12 months with any person described in items 1-4 or with a person known or suspected to have HIV infection	Women who have had sex with a man with a history of MSM behavior in the preceding 12 months
#4- Men and women who have engaged in sex in exchange for money or drugs in the preceding 5 years	People who have had sex in exchange for money or drugs in the preceding 12 months
#5- Persons who have had sex in the preceding 12 months with any person described in items 1-4 or with a person known or suspected to have HIV infection	People who have had sex with a person who had sex in exchange for money or drugs in the preceding 12 months
#5- Persons who have had sex in the preceding 12 months with any person described in items 1-4 or with a person known or suspected to have HIV infection	People who have had sex with a person that has injected drugs by IV, IM, or subQ route for nonmedical reasons in the preceding 12 months
<p><i>Specific Exclusionary Criteria for Ped Donors #2-</i> Children born to mothers with HIV infection or mothers who meet the behavioral or laboratory exclusionary criteria for adult donors (regardless of HIV status) should not be accepted as donors unless HIV infection can be definitely excluded in the child as follows:</p> <p>Children >18 months of age who are born to mothers with or at risk for HIV infection, who have not been breastfed within the last 12 months, and whose HIV Ab tests, physical exam, and review of medical records don not indicate evidence of HIV infection can be accepted as donors</p>	A child who is ≤ 18 months of age and born to a mother known to be infected with, or at increased risk for HIV, HBV, or HCV infections
<p><i>Specific Exclusionary Criteria for Ped Donors #3-</i> Children ≤ 18 months of age who are born to mothers with or at risk for HIV infection or who have been breastfed within the past 12 months should not be accepted as donors regardless of their HIV status</p>	A child who has been breastfed within the preceding 12 months and the mother is known to be infected with, or at increased risk for HIV infection (<i>Recommendation 4</i>)
#2- Persons who report nonmedical IV, IM, or subQ injection of drugs in the preceding 5 years	People who have injected drugs by IV, IM, or subQ route for nonmedical reasons in the preceding 12 months

#7- Inmates of correctional systems. (This exclusion is to address issues such as difficulties with informed consent and increased prevalence of HIV in this population.)	People who have been in lockup, jail, prison, or a juvenile correctional facility for more than 72 hours in the preceding 12 months
<i>Laboratory and Other Medical Exclusionary Criteria</i> #3- Persons whose history, physical exam, medical records, or autopsy reports reveal other evidence of HIV infection or high-risk behavior, such as a diagnosis of AIDS, unexplained weight loss, night sweats, blue or purple spots on the skin or mucous membranes typical of Kaposi's Sarcoma, unexplained lymphadenopathy lasting ≥ 1 month, unexplained temperature >100.5 F (38.6 C) for >10 days, unexplained persistent cough, MSM contact, sexually transmitted diseases , or needle tracks or other signs of parenteral drug abuse.	People who have been newly diagnosed with or have been treated for syphilis, gonorrhea, <i>Chlamydia</i> , or genital ulcers in the preceding 12 months
<i>NEW</i>	<i>Donors who meet the following criterion should be identified as being at increased risk for recent HCV infection only:</i> People who have been on hemodialysis in the preceding 12 months
#3-Persons with hemophilia or related clotting disorders who have received human-derived clotting factor concentrates	<i>Not addressed specifically in 2013 med-soc criteria</i>
#6- Person who have been exposed in the preceding 12 months to known or suspected HIV-infected blood through percutaneous inoculation or through contact with an open wound, nonintact skin, or mucous membrane.	<i>Not addressed specifically in 2013 med-soc criteria</i>
<i>Laboratory and Other Medical Exclusionary Criteria</i>	
#1- Persons who cannot be tested for HIV infection because of refusal, inadequate blood samples (e.g. hemodilution that could result in a false negative test), or any other reasons	<i>Hemodilution specifically addressed in Recommendation 5b</i>
#2- Persons with a repeatedly reactive screening assay for HIV-1 or HIV-2 Ab regardless of the result of supplemental assays.	